

Spice oleoresins 30 parts per million (provided that if residues of other chlorinated solvents are also present, the total of all residues of such solvents in spice oleoresins shall not exceed 30 parts per million).

Subpart D—Specific Usage Additives

§ 173.300 Chlorine dioxide.

Chlorine dioxide (CAS Reg. No. 10049-04-4) may be safely used in food in accordance with the following prescribed conditions:

(a) The additive is generated by treating an aqueous solution of sodium chlorite with either chlorine gas or a mixture of sodium hypochlorite and hydrochloric acid. The generator effluent contains at least 90 percent (by weight) of chlorine dioxide with respect to all chlorine species as determined by Method 4500-ClO₂ E in the "Standard Methods for the Examination of Water and Wastewater," 18th ed., 1992, or an equivalent method. Method 4500-ClO₂ E is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the Center for Food Safety and Applied Nutrition (HFS-200), Food And Drug Administration, 200 C St., SW., Washington, DC 20204-0001 and The American Public Health Association, 1015 Fifteenth St., NW., Washington, DC 20005, or may be examined at the Office of the Federal Register, 800 North Capitol St., NW., suite 700, Washington, DC.

(b)(1) The additive may be used as an antimicrobial agent in water used in poultry processing in an amount not to exceed 3 parts per million (ppm) residual chlorine dioxide as determined by Method 4500-ClO₂ E, referenced in paragraph (a) of this section, or an equivalent method.

(2) The additive may be used as an antimicrobial agent in water used to wash fruits and vegetables that are not raw agricultural commodities in an amount not to exceed 3 ppm residual chlorine dioxide as determined by Method 4500-ClO₂ E, referenced in paragraph (a) of this section, or an equivalent method. Treatment of the fruits and vegetables with chlorine dioxide shall be followed by a potable water rinse or by blanching, cooking, or canning.

[60 FR 11900, Mar. 3, 1995. Redesignated at 61 FR 14245, Apr. 1, 1996, as amended at 61 FR 14480, Apr. 2, 1996; 63 FR 38747, July 20, 1998]

§ 173.310 Boiler water additives.

Boiler water additives may be safely used in the preparation of steam that will contact food, under the following conditions:

(a) The amount of additive is not in excess of that required for its functional purpose, and the amount of steam in contact with food does not exceed that required to produce the intended effect in or on the food.

(b) The compounds are prepared from substances identified in paragraphs (c) and (d) of this section, and are subject to the limitations, if any, prescribed:

(c) List of substances:

Substances	Limitations
Acrylamide-sodium acrylate resin	Contains not more than 0.05 percent by weight of acrylamide monomer.
Acrylic acid/2-acrylamido-2-methyl propane sulfonic acid copolymer having a minimum weight average molecular weight of 9,900 and a minimum number average molecular weight of 5,700 as determined by a method entitled "Determination of Weight Average and Number Average Molecular Weight of 60/40 AA/AMPS" (October 23, 1987), which is incorporated by reference in accordance with 5 U.S.C. 552(a). Copies may be obtained from the Center for Food Safety and Applied Nutrition (HFS-200), Food and Drug Administration, 200 C St. SW., Washington, DC 20204, or may be examined at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC.	Total not to exceed 20 parts per million (active) in boiler feedwater.
Ammonium alginate.	
Cobalt sulfate (as catalyst).	
1-hydroxyethylidene-1,1-diphosphonic acid (CAS Reg. No. 2809-21-4) and its sodium and potassium salts.	
Lignosulfonic acid.	

Substances	Limitations
Monobutyl ethers of polyethylene-polypropylene glycol produced by random condensation of a 1:1 mixture by weight of ethylene oxide and propylene oxide with butanol.	Minimum mol. wt. 1,500.
Poly(acrylic acid-co-hypophosphite), sodium salt [CAS Reg. No. 71050–62–9], produced from a 4:1 to a 16:1 mixture by weight of acrylic acid and sodium hypophosphite.	Total not to exceed 1.5 parts per million in boiler feed water. Copolymer contains not more than 0.5 percent by weight of acrylic acid monomer (dry weight basis). As defined in § 172.820 of this chapter.
Polyethylene glycol	Total not to exceed 1 part per million in boiler feed water (calculated as the acid).
Polymaleic acid [CAS Reg. No. 26099–09–2], and/or its sodium salt. [CAS Reg. No. 30915–61–8 or CAS Reg. No. 70247–90–4].	Minimum mol. wt. 1,000.
Polyoxypropylene glycol	
Potassium carbonate.	
Potassium tripolyphosphate.	
Sodium acetate.	
Sodium alginate.	
Sodium aluminate.	
Sodium carbonate.	
Sodium carboxymethylcellulose	Contains not less than 95 percent sodium carboxymethylcellulose on a dry-weight basis, with maximum substitution of 0.9 carboxymethylcellulose groups per anhydroglucose unit, and with a minimum viscosity of 15 centipoises for 2 percent by weight aqueous solution at 25 °C; by the method prescribed in the "Food Chemicals Codex," 4th ed. (1996), pp. 744–745, which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the National Academy Press, Box 285, 2101 Constitution Ave. NW., Washington, DC 20055 (Internet address "http://www.nap.edu"), or may be examined at the Center for Food Safety and Applied Nutrition's Library, Food and Drug Administration, 200 C St. SW., rm. 3321, Washington, DC, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.
Sodium glucoheptonate	Less than 1 part per million cyanide in the sodium glucoheptonate.
Sodium hexametaphosphate.	
Sodium humate.	
Sodium hydroxide.	
Sodium lignosulfonate.	
Sodium metabisulfite.	
Sodium metasilicate.	
Sodium nitrate.	
Sodium phosphate (mono-, di-, tri-).	
Sodium polyacrylate.	
Sodium polymethacrylate.	
Sodium silicate.	
Sodium sulfate.	
Sodium sulfite (neutral or alkaline).	
Sodium tripolyphosphate.	
Tannin (including quebracho extract).	
Tetrasodium EDTA.	
Tetrasodium pyrophosphate.	

(d) Substances used alone or in combination with substances in paragraph (c) of this section:

Substances	Limitations
Cyclohexylamine	Not to exceed 10 parts per million in steam, and excluding use of such steam in contact with milk and milk products.
Diethylaminoethanol	Not to exceed 15 parts per million in steam, and excluding use of such steam in contact with milk and milk products.
Hydrazine	Zero in steam.
Morpholine	Not to exceed 10 parts per million in steam, and excluding use of such steam in contact with milk and milk products.
Octadecylamine	Not to exceed 3 parts per million in steam, and excluding use of such steam in contact with milk and milk products.
Trisodium nitrilotriacetate	Not to exceed 5 parts per million in boiler feedwater; not to be used where steam will be in contact with milk and milk products.

(e) To assure safe use of the additive, in addition to the other information required by the Act, the label or labeling shall bear:

(1) The common or chemical name or names of the additive or additives.

(2) Adequate directions for use to assure compliance with all the provisions of this section.

[42 FR 14526, Mar. 15, 1977, as amended at 45 FR 73922, Nov. 7, 1980; 45 FR 85726, Dec. 30, 1980; 48 FR 7439, Feb. 22, 1983; 49 FR 5748, Feb. 15, 1984; 49 FR 10106, Mar. 19, 1984; 50 FR 49536, Dec. 3, 1985; 53 FR 15199, Apr. 28, 1988; 54 FR 31012, July 26, 1989; 55 FR 12172, Apr. 2, 1990; 61 FR 14245, Apr. 1, 1996; 64 FR 1759, Jan. 12, 1999]

§ 173.315 Chemicals used in washing or to assist in the peeling of fruits and vegetables.

Chemicals may be safely used to wash or to assist in the peeling of fruits and vegetables in accordance with the following conditions:

(a) The chemicals consist of one or more of the following:

(1) Substances generally recognized as safe in food or covered by prior sanctions for use in washing fruits and vegetables.

(2) Substances identified in this subparagraph and subject to such limitations as are provided:

Substances	Limitations
A mixture of alkylene oxide adducts of alkyl alcohols and phosphate esters of alkylene oxide adducts of alkyl alcohols consisting of: α -alkyl (C_{12} - C_{18})- <i>omega</i> -hydroxy-poly (oxyethylene) (7.5–8.5 moles)/poly (oxypropylene) block copolymer having an average molecular weight of 810; α -alkyl (C_{12} - C_{18})- <i>omega</i> -hydroxy-poly (oxyethylene) (3.3–3.7 moles) polymer having an average molecular weight of 380, and subsequently esterified with 1.25 moles phosphoric anhydride; and α -alkyl (C_{10} - C_{12})- <i>omega</i> -hydroxypoly (oxyethylene) (11.9–12.9 moles)/poly (oxypropylene) copolymer, having an average molecular weight of 810, and subsequently esterified with 1.25 moles phosphoric anhydride.	May be used at a level not to exceed 0.2 percent in lye-peeling solution to assist in the lye peeling of fruit and vegetables.
Aliphatic acid mixture consisting of valeric, caproic, enanthic, caprylic, and pelargonic acids.	May be used at a level not to exceed 1 percent in lye peeling solution to assist in the lye peeling of fruits and vegetables.
1-Hydroxyethylidene-1,1-diphosphonic acid	May be used only with peroxyacetic acid. Not to exceed 4.8 ppm in wash water. Limited to use on fruits and vegetables that are not raw agricultural commodities.
Hydrogen peroxide	Used in combination with acetic acid to form peroxyacetic acid. Not to exceed 59 ppm in wash water. Limited to use on fruits and vegetables that are not raw agricultural commodities.
Peroxyacetic acid	Prepared by reacting acetic acid with hydrogen peroxide. Not to exceed 80 ppm in wash water. Limited to use on fruits and vegetables that are not raw agricultural commodities.
Polyacrylamide	Not to exceed 10 parts per million in wash water. Contains not more than 0.2 percent acrylamide monomer. May be used in the washing of fruits and vegetables.
Potassium bromide	May be used in the washing or to assist in the lye peeling of fruits and vegetables.
Sodium <i>n</i> -alkylbenzene-sulfonate (alkyl group predominantly C_{12} and C_{13} and not less than 95 percent C_{10} to C_{16}).	Not to exceed 0.2 percent in wash water. May be used in washing or to assist in the lye peeling of fruits and vegetables.
Sodium dodecylbenzene-sulfonate (alkyl group predominantly C_{12} and not less than 95% C_{10} to C_{16}).	Do.
Sodium 2 ethyl-hexyl sulfate	Do.
Sodium hypochlorite	May be used in the washing or to assist in the lye peeling of fruits and vegetables.
Sodium mono- and dimethyl naphthalene sulfonates (mol. wt. 245–260)	Not to exceed 0.2 percent in wash water. May be used in the washing or to assist in the lye peeling of fruits and vegetables.

(3) Sodium mono- and dimethyl naphthalene sulfonates (mol. wt. 245–260) may be used in the steam/scald vacuum peeling of tomatoes at a level not to

exceed 0.2 percent in the condensate or scald water.

(4) Substances identified in this paragraph (a)(4) for use in flume water for washing sugar beets prior to the slicing